


SPOZNAJTE OPIČIE KIAHNE



ÚRAD
VEREJNÉHO
ZDRAVOTNÍCTVA
SLOVENSKEJ
REPUBLIKY

Opičie kiahne sa môžu prenášať blízky fyzickým kontaktom (koža na kožu), a to aj prostredníctvom intímneho kontaktu alebo počas sexuálneho styku.



AKO SA OCHORENIE ŠÍRI



Úzkym kontaktom s:



vyrážkami ako aj výlučkami z nich



kontaminovanými predmetmi

ide o predmety ako napr. uteráky, oblečenie, riady, posteľná bielizeň, sexuálne hračky



telesnými tekutinami alebo veľkými kvapôčkami priamo z tváre do tváre


je potrebný dlhší kontakt než v prípade COVID-19

! Inkubačný čas môže byť 5 až 21 dní

AKÉ SÚ PRÍZNAKY OCHORENIA



Horúčka alebo zimnica



Opuch lymfatických uzlín



Silná bolesť hlavy



Bolesti svalov či chrbta



Únava alebo vyčerpanie



Vyrážka sa objaví zvyčajne 1 až 3 dni od výskytu niektorého z príznakov

Pľuzgieriky sa môžu objaviť aj na dlaniach, chodidlách, vnútri úst, pri análnom otvore alebo na genitáliách. Môže ísť aj o ojedinelý pľuzgierik.

Infekcia sa môže ďalej prenášať až do zahojenia poslednej chrasty **!**


Pri podozrení na ochorenie alebo úzky kontakt s chorým konzultujte vhodný postup s lekárom. Ak je potrebné vyšetrenie, vyrážku si mimo domácnosti prekryte oblečením a nasadte si respirátor.

RECOGNIZING MONKEYPOX



PUBLIC
HEALTH
AUTHORITY
OF THE SLOVAK
REPUBLIC

Monkeypox can be transmitted through close physical contact (skin-to-skin), including intimate or sexual contact.



HOW THE DISEASE SPREADS

Close contact with:




! The incubation period ranges from 5 to 21 days



contaminated objects

items such as towels, clothes, dishes, bed linen, sex toys, etc.



rash, including secretions from pox lesions


body fluids or large droplets during close (face-to-face) contact

longer contact is needed compared to COVID-19

WHAT ARE THE SYMPTOMS




Fever or chills



Swollen lymph nodes



Strong headache



Muscle aches or backache



Fatigue or exhaustion

The rash (pimples or blisters) typically appears 1 to 3 days after initial symptoms.



Blisters can also appear on the palms of the hands, feet soles, inside the mouth, near the anus or on the genitals. It can also be a single blister.

! Infection can be transmitted until the last scab heals

If you have symptoms or if you had close contact with the infected person, you should talk to your healthcare provider. If personal examination is necessary, cover the rash with clothing and wear a respirator when leaving home.